

2020-08

# A novel tri-unsaturated highly branched isoprenoid (HBI) alkene from the marine diatom *Navicula salinicola*

Gao, S

<http://hdl.handle.net/10026.1/15905>

---

10.1016/j.orggeochem.2020.104050

Organic Geochemistry

Elsevier BV

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*

**A novel tri-unsaturated highly branched isoprenoid (HBI) alkene  
from the marine diatom *Navicula salinicola***

Simin Gao <sup>a,b,\*</sup>, Lukas Smik <sup>c</sup>, Maxim Kulikovskiy <sup>d</sup>, Nataliya Shkurina <sup>d</sup>,  
Evgeniy Gusev<sup>d</sup>, Nikolai Pedentchouk <sup>a,d</sup>, Thomas Mock <sup>a</sup>, Simon T. Belt <sup>c</sup>

(a) *School of Environmental Sciences, University of East Anglia, Norwich*

*Research Park, Norwich, NR4 7TJ, UK.*

(b) *Department of Ocean Science and Engineering, Southern University of*

*Science and Technology, Shenzhen, China.*

(c) *School of Geography, Earth and Environmental Sciences, University*

*of Plymouth, Drake Circus, Plymouth, Devon PL4 8AA, UK.*

(d) *K.A. Timiryazev Institute of Plant Physiology RAS, Moscow, Russia*

\* Corresponding author, [simin.gao@uea.ac.uk](mailto:simin.gao@uea.ac.uk)

**ABSTRACT**

A novel tri-unsaturated C<sub>25</sub> highly branched isoprenoid (HBI) alkene has been identified in a laboratory culture of the diatom *Navicula salinicola* and its structure determined using a combination of NMR spectroscopy and gas chromatography–mass spectrometry (GC–MS). This represents the first report of a C<sub>25</sub> HBI in a marine diatom from the *Navicula* genus, although a different tri-unsaturated C<sub>25</sub> HBI has been reported previously in the freshwater species *N. sclesvicensis* and unspecified HBIs have been identified in the brackish *N. phyllepta*. The newly characterised HBI contains a relatively unusual conjugated diene sub-unit, a structural feature

only previously reported in some HBIs biosynthesised by a further marine diatom, *Haslea ostrearia*.

**Keywords:** highly branched isoprenoid; alkene; diatom; *Navicula salinicola*

## 1. Introduction

C<sub>25</sub> highly branched isoprenoid (HBI) alkenes are common components of marine and lacustrine sediments worldwide and are biosynthesised by certain diatoms mainly belonging to the genera *Haslea*, *Navicula*, *Rhizosolenia*, *Pleurosigma*, *Berkeleya* and *Pseudosolenia* (Volkman et al., 1994; Belt et al., 1996, 2000, 2001; Sinninghe Damsté et al., 1999; Grossi et al., 2004; Brown et al., 2014; Kaiser et al., 2016). A single tri-unsaturated C<sub>25</sub> HBI (Structure 6; Fig. 1) has been identified in the freshwater diatom *Navicula sclesvicensis* (Belt et al., 2001), and the mainly brackish species *N. phyllepta* has also been reported as an HBI-producing diatom (Sinninghe Damsté et al., 2004), although no structures were given. In contrast, no HBIs have as yet been reported in any marine species within the *Navicula* genus. Here, we identify a novel tri-unsaturated C<sub>25</sub> HBI isolated from a laboratory culture of the marine diatom *N. salinicola* and report its structure based on analysis by NMR spectroscopy and gas chromatography–mass spectrometry (GC–MS).

## 2. Experimental

The benthic diatom *N. salinicola* was collected from a coastal marine environment (M. Kulikovskiy, June 2016 at Nha Trang, Vietnam, 12°13'14.5"N 109°12'18.3"E) and kept as strain BTD1 in the Laboratory of

46 Molecular Taxonomy of Aquatic Plants Institute of Plant Physiology (RAS).  
47 (Further taxonomic information can be found in the Supplementary  
48 Information). Initially, *N. salinicola* was cultured at 15 °C, 150  $\mu\text{mol m}^{-2} \text{s}^{-1}$   
49 continuous light in small flasks (150ml). Samples were harvested by  
50 filtration using MF-Millipore™ membrane filters (25 mm diameter, 0.3  $\mu\text{m}$   
51 pore size) during the exponential and stationary growth phases (Fig. 2).  
52 Large-scale cultures were then set up in several 2 l conical flasks under the  
53 same growth conditions. 80 l of such cultures were harvested by  
54 centrifugation (4000 rpm, 10 mins) during the stationary growth phase. For  
55 the small- and large-scale culture experiments, we used enriched f/2  
56 medium (Guillard and Ryther 1962), along with the following nutrient  
57 concentrations: 2646  $\mu\text{M}$   $\text{NaNO}_3$ , 318  $\mu\text{M}$   $\text{Na}_2\text{SiO}_3$ , 108  $\mu\text{M}$   $\text{NaH}_2\text{PO}_4$ . The  
58 filtered and centrifuged biomass was freeze dried and the resulting dry  
59 material extracted via sonication using hexane (3 ml (small-scale cultures);  
60 25 ml (large-scale culture)). The total hexane extract (THE) was then  
61 concentrated by removing hexane under a stream of nitrogen and partially  
62 purified using column chromatography ( $\text{SiO}_2$ ). The hydrocarbon fraction  
63 (hexane) was analysed by GC–MS using an Agilent 7890 gas chromatograph  
64 equipped with a HP5MS fused-silica column (30 m; 0.25  $\mu\text{m}$  film thickness;  
65 0.25 mm internal diameter) coupled to an Agilent 5975 series Mass  
66 Selective Detector (MSD). NMR data were obtained using a JEOL ECP-400  
67 NMR spectrometer with chemical shifts measured relative to those of  $\text{CDCl}_3$   
68 ( $^1\text{H}$ : 7.24 ppm;  $^{13}\text{C}$ : 77.0 ppm). NMR data were collected on the THE

obtained from the large-scale culture. The purity of the newly reported C<sub>25:3</sub> HBI (see Section 3) in this THE was estimated to be ca. 90% based on its relative peak area (GC–MS; Supplementary Information) and by the relative integration values of H-23 (Fig. 1) versus the alkenic protons of the co-occurring polyunsaturated linear alkenes ( $\delta$  = ca. 5.3 ppm) in the <sup>1</sup>H NMR spectrum. ).

### 3. Results and discussion

Following extraction of several small-scale cultures of *N. salinicola* from the exponential and stationary growth phases, analysis of partially purified THEs by GC–MS revealed the presence of a suite of closely eluting polyunsaturated linear alkenes (e.g. heneicosa-3,6,9,12,15,18-hexaene; *n*-C<sub>21:6</sub>), trace amounts of a di-unsaturated HBI (**2**; Fig. 1) and a further compound exhibiting similar mass spectral properties to a range of C<sub>25</sub> HBIs characterised previously. However, although the retention index (RI) of this component (RI<sub>HP5ms</sub> = 2141) did not match that of any previously reported C<sub>25</sub> HBIs, hydrogenation of an aliquot of one THE resulted in the formation of the parent HBI alkane C<sub>25:0</sub>, thus confirming the C<sub>25</sub> carbon skeleton. Interestingly, this new HBI was only detected in cultures harvested during the stationary phase and was identified as tri-unsaturated on the basis of its molecular ion (M<sup>+</sup>. 346; Fig. 3). At this point, it is not clear why this new HBI and the co-occurring diene **2** were not detected during the exponential growth phase, although we note that some variability in cellular HBI concentrations and distributions have been reported in a small number of

previous studies (e.g. Wraige et al., 1997,1998,1999; Brown et al., 2020). From the large-scale culture of *N. salinicola*, we obtained ca 0.2 mg of the partially purified HBI triene (**3**; Fig. 1), which enabled full structural characterisation using  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectroscopy. A conjugated diene sub-structure (C22–C25; C23–C24; Fig 1) could be readily identified through a particularly characteristic low field resonance in the  $^1\text{H}$  NMR spectrum due to H-23 (Wraige et al., 1997; Allard et al., 2001), together with further low field resonances that could be attributed to alkenic methylene protons at C24 and C25 (Table 1). The third double bond could also be identified from its methylene protons at C17. Alternative positions for this third double bond at C1–C2 or C14–C15 can be discounted due to the observation of two isopropyl groups in the  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra (Table 1). Further, a double bond at C10–C18 leaves a solitary methyl group at C17 whose  $^{13}\text{C}$  chemical shift would be at ca. 15.5 ppm, by comparison with related compounds (e.g. **1**; Belt et al., 2012). In contrast, isolated methyl groups at C18 in previously characterised HBIs resonate at ca. 19–20 ppm (e.g. 19.8 ppm for HBIs **1** and **2**; Fig. 1) (Johns et al., 1999; Belt et al., 2012), consistent with that observed for HBI **3** (i.e. 19.9 ppm; Table 1). The  $^{13}\text{C}$  NMR spectrum of HBI **3** also contained individual resonances due to the six magnetically inequivalent alkenic carbon nuclei (Table 1) and complete  $^{13}\text{C}$  resonance assignments could be proposed by comparison with structurally similar HBIs characterised previously (Fig. 1; Belt et al., 1996, 2012; Wraige et al., 1997; Allard et al., 2001), some of which contain a conjugated diene

sub-unit (i.e. **7–8**). The GC RI of HBI **3** ( $RI_{HP5ms} = 2141$ ) is substantially higher than those of some other HBI trienes (e.g.  $RI_{HP5ms} = 2114$  (**4**); 2109 (**5**); 2090 (**6**)) but is consistent with that reported for the structurally related tetraene **7** ( $RI_{HP-5} = 2159$ ; Allard et al., 2001).

The identification of a  $C_{25}$  HBI in *N. salinicola* represents the first example of HBI production within a marine *Navicula* species despite the near-ubiquity of this genus within natural diatom populations. Since the *Navicula* and *Haslea* genera are quite similar, phylogenetically, with the latter well-known as an HBI-producing genus, the new finding is probably not surprising; however *Navicula* is a far more common genus, potentially making it a more important source of some HBIs in marine sediments. In terms of its structure, the conjugated diene sub-structure (C22–C25; C23–C24; Fig. 1) is somewhat unusual, although there is some previous precedent for such a feature in other HBIs isolated from a small number of cultures of the marine diatom *Haslea ostrearia* (Wraige et al., 1997; Allard et al., 2001). In fact, HBI **3** is a close structural analogue of HBI **7** identified previously in *H. ostrearia*, albeit as a minor component. Since the retention index of HBI **3** ( $RI_{HP5ms} = 2141$ ) is similar to two HBI tetraenes identified in some cultures of *H. ostrearia* (viz.  $RI_{HP-5} = 2143–2146$ ; Allard et al., 2001), one of which also contained HBI **7**, it is possible that HBI **3** may also have been present in the corresponding lipid extracts, but not identified due to co-elution. We are unaware of any geochemical reports of HBI **3** although its characterisation described herein may, in the future, lead to its positive

identification in sedimentary archives, an outcome that may potentially add to the use of HBIs as palaeoenvironmental indicators (c.f. HBIs 1 and 2 for Arctic and Antarctic sea ice; see Belt, 2018 for a review).

#### 4. Conclusions

We report the structural identification of a novel C<sub>25</sub> HBI biomarker in the marine diatom *N. salinicola*, the first example of HBI production within a marine *Navicula* species, thus expanding the potential number of sources of HBIs in the environment. Further studies into *N. salinicola* and related species may prove valuable in the use of HBIs as palaeoenvironmental proxies.

#### Acknowledgments

SG was funded by the UEA-SUSTech Split-Site PhD programme. We also acknowledge partial support by The Leverhulme Trust (RPG-2019-088). MK, EG, and NS were supported by Russian Science Foundation (19-14-00320) and by The Framework of The State Assignment (theme AAAA-A19-119041190086-6). We also thank two anonymous reviewers who provided helpful feedback on the original manuscript.

#### References

- Allard, W.G., Belt, S.T., Massé, G., Naumann, R., Robert, J.-M., Rowland, S.J., 2001. Tetra-unsaturated sesterterpenoids (Haslenes) from *Haslea ostrearia* and related species. *Phytochemistry* 56, 795–800.
- Belt, S.T., Cooke, D.A., Robert, J.-M., Rowland, S.J., 1996. Structural characterisation of widespread polyunsaturated isoprenoid biomarkers:



- 161 A C<sub>25</sub> triene, tetraene and pentaene from the diatom *Haslea ostrearia*  
162 Simonsen. Tetrahedron Letters 37, 4755–4758.
- 163 Belt, S.T., Allard, W.G., Massé, G., Robert, J.-M., Rowland, S.J., 2000.  
164 Highly branched isoprenoids (HBIs): identification of the most  
165 common and abundant sedimentary isomers. Geochimica et  
166 Cosmochimica Acta 64, 3839–3851.
- 167 Belt, S.T., Massé, G., Allard, W.G., Robert, J.-M., Rowland, S.J., 2001.  
168 Identification of a C<sub>25</sub> highly branched isoprenoid triene in the  
169 freshwater diatom *Navicula sclesvicensis*. Organic Geochemistry 32,  
170 1169-1172.
- 171 Belt, S.T., Brown, T.A., Cabedo-Sanz, P., Navarro-Rodriguez, A., 2012.  
172 Structural confirmation of the sea ice biomarker IP<sub>25</sub> found in Arctic  
173 marine sediments. Environmental Chemistry Letters 10, 189–192.
- 174 Brown, T.A., Belt, S.T., Cabedo-Sanz, P., 2014. Identification of a novel di-  
175 unsaturated C<sub>25</sub> highly branched isoprenoid in the marine tube-  
176 dwelling diatom *Berkeleya rutilans*. Environmental Chemistry  
177 Letters, 12, 455–460.
- 178 Brown, T.A., Rad-Menéndez, C., Ray, J.L., Skaar, K.S., Thomas, N., Ruiz-  
179 Gonzalez, C., Leu, E., 2020. Influence of nutrient availability on  
180 Arctic sea ice diatom HBI lipid synthesis. Organic Geochemistry 141,  
181 103977.
- 182 Grossi, V., Beker, B., Geenevasen, J.A.J., Schouten, S., Raphel, D., Fontaine,  
183 M.-F., Sinninghe Damsté, J.S., 2004. C<sub>25</sub> highly branched isoprenoid

184 alkenes from the marine benthic diatom *Pleurosigma strigosum*.  
 185 Phytochemistry 65, 3049–3055.

186 Guillard, R.R. L., and Ryther, J.H., 1962. Studies of marine planktonic  
 187 diatoms. I. *Cyclotella nana* Hustedt and *Detonula confervacea* Cleve.  
 188 Canadian Journal of Microbiology 8, 229–239.

189 Johns, L., Wraige, E.J., Belt, S.T., Lewis, C.A., Massé, G., Robert, J.-M.,  
 190 Rowland, S.J., 1999. Identification of a C<sub>25</sub> highly branched isoprenoid  
 191 (HBI) diene in Antarctic sediments, Antarctic sea-ice diatoms and  
 192 cultured diatoms. Organic Geochemistry 30, 1471–1475.

193 Kaiser, J., Belt, S.T., Tomczak, T., Brown, T.A., Wasmund, N., Arz, H.W.,  
 194 2016. C<sub>25</sub> highly branched isoprenoid alkenes in the Baltic Sea  
 195 produced by the marine planktonic diatom *Pseudosolenia calcar-avis*.  
 196 Organic Geochemistry 93, 51–58.

197 Sinninghe Damsté, J.S., Schouten, S., Rijpstra, W.I.C., Hopmans, E.C.,  
 198 Peletier, H., Gieskes, W.W.C., Geenevasen, J.A.J., 1999. Structural  
 199 identification of the C<sub>25</sub> highly branched isoprenoid pentaene in the  
 200 marine diatom *Rhizosolenia setigera*. Organic Geochemistry 30, 1581-  
 201 1583.

202 Sinninghe Damsté, J.S., Muyzer, G., Abbas, B., Rampen, S.W., Massé, G.,  
 203 Allard, W.G., Belt, S.T., Robert, J.-M., Rowland, S.J., Moldowan, J.M.,  
 204 Barbanti, S.M., Fago, F.J., Denisevich, P., Dahl, J., Trindade, L.A.F.,  
 205 Schouten, S., 2004. The rise of the rhizosolenid diatoms. Science 304,  
 206 584–587.

207 Volkman, J.K., Barrett, S.M., Dunstan, G.A., 1994. C<sub>25</sub> and C<sub>30</sub> highly  
 208 branched isoprenoid alkenes in laboratory cultures of two marine  
 209 diatoms. *Organic Geochemistry* 21, 407–414.

210 Wraige, E.J., Belt, S.T., Lewis, C.A., Cooke, D.A., Robert, J.-M., Massé, G.,  
 211 Rowland, S.J., 1997. Variations in structures and distributions of C<sub>25</sub>  
 212 highly branched isoprenoid (HBI) alkenes in cultures of the diatom,  
 213 *Haslea ostrearia* (Simonsen). *Organic Geochemistry* 27, 497–505.

214 Wraige, E.J., Belt, S.T., Massé, G., Robert, J.-M., Rowland, S.J., 1998.  
 215 Variations in distributions of C<sub>25</sub> highly branched isoprenoid (HBI)  
 216 alkenes in the diatom, *Haslea ostrearia*: influence of salinity. *Organic*  
 217 *Geochemistry* 28, 855–859.

218 Wraige, E.J., Johns, L., Belt, S.T., Massé, G., Robert, J.-M., Rowland, S.J.,  
 219 1999. Highly branched C<sub>25</sub> isoprenoids in axenic cultures of *Haslea*  
 220 *ostrearia*. *Phytochemistry* 51, 69–73.

## 221 **Figure legends**

222 **Figure 1.** Structures of C<sub>25</sub> HBIs referred to in the text. HBIs are numbered  
 223 in order of increasing unsaturation.

224 **Figure 2.** Growth curve of the small-scale culture of the marine diatom *N.*  
 225 *salinicola*. Cell densities were estimated by measuring in vivo fluorescence  
 226 between 460 nm to 670 nm using the SpectraMax® iD3 Multi-Mode  
 227 Microplate Reader. Cultures were harvested during the exponential (empty  
 228 arrow) and/or stationary (solid arrow) growth phases (n=1).

229 **Figure 3.** Structure and mass spectrum of HBI 3.

Figure 1.




Figure 2.





Figure 3.



249

250 Table 1. Key NMR data for HBI 3.

|        | Carbon shift ( $\delta$ /ppm) | Proton Number | Proton shift ( $\delta$ /ppm)    |
|--------|-------------------------------|---------------|----------------------------------|
| 1,16*  | 22.8, 22.7*                   | 1,15,16,19    | 0.85 (12H, m)                    |
| 2      | 28.0                          | 5,7,21        | 2.05 (5H, m)                     |
| 3      | 39.1                          | 17            | 4.77, 4.72 (2H, 2 x s, br)       |
| 4      | 25.6                          | 18            | 0.82 (3H, t, $J=6.9$ Hz)         |
| 5      | 33.0                          | 23            | 6.33 (1H, dd, $J=17.6, 11.0$ Hz) |
| 6      | 152.3                         | 24a           | 5.01 (1H, d, $J=11.0$ Hz)        |
| 7      | 47.1                          | 24b           | 5.16 (1H, d, $J=17.6$ Hz)        |
| 8      | 29.8                          | 25            | 4.96 (2H, m, br)                 |
| 9      | 34.9                          |               |                                  |
| 10     | 33.0                          |               |                                  |
| 11     | 37.1                          |               |                                  |
| 12     | 24.8                          |               |                                  |
| 13     | 39.4                          |               |                                  |
| 14     | 28.0                          |               |                                  |
| 15,19* | 22.8, 22.7*                   |               |                                  |
| 17     | 109.0                         |               |                                  |
| 18     | 19.9                          |               |                                  |
| 20     | 29.4                          |               |                                  |
| 21     | 32.0                          |               |                                  |
| 22     | 146.9                         |               |                                  |
| 23     | 139.0                         |               |                                  |
| 24     | 113.1                         |               |                                  |
| 25     | 115.5                         |               |                                  |

251

\*Assignments may be interchanged